



Contents lists available at ScienceDirect

Journal of Clinical Neuroscience

journal homepage: www.elsevier.com/locate/jocn

Pitfalls in the diagnosis of pupil-sparing oculomotor nerve palsy without limb ataxia: A case report of a variant of Claude's syndrome and neuroanatomical analysis using diffusion-tensor imaging

Eiichiro Amano^{a,*}, Tetsuya Komatuzaki^b, Hideaki Ishido^a, Tasuku Ishihara^a, Shinichi Otsu^a, Ichiro Yamada^c, Akira Machida^a

^a Tsuchiura Kyodo General Hospital, Department of Neurology, 4-1-1 Otsuno, Tsuchiura-shi, Ibaraki 300-0028, Japan

^b Tsuchiura Kyodo General Hospital, Department of Radiology, 4-1-1 Otsuno, Tsuchiura-shi, Ibaraki 300-0028, Japan

^c Tokyo Medical and Dental University, Department of Radiology, 1-5-45 Yushima, Bunkyo-ku, Tokyo 113-8510, Japan

ARTICLE INFO

Article history:

Received 30 August 2017

Accepted 30 September 2017

Available online xxx

Keywords:

Oculomotor nerve palsy

Truncal ataxia

Magnetic resonance imaging

Superior cerebellar peduncle

Tectospinal tract

ABSTRACT

Midbrain infarction causing oculomotor nerve palsy with contralateral ataxia is named Claude's syndrome. Herein we report the case of a variant of Claude's syndrome, which shows pupil-sparing oculomotor nerve palsy without the accompanying neurological deficits other than subtle truncal ataxia. MRI and Diffusion Tensor Imaging revealed that midbrain infarction was located rostrally above the decussation of the superior cerebellar peduncle (SCP) and might have partially destructed the tectospinal tract, which resulted in the absence of limb ataxia and presence of subtle truncal ataxia. In this variant of Claude's syndrome, we should carefully assess truncal ataxia to avoid misdiagnosing it as isolated pupil-sparing oculomotor nerve palsy because the patient showed apparently normal gait and truncal ataxia was only revealed by unstable tandem gait.

© 2017 Elsevier Ltd. All rights reserved.

1. Introduction

Isolated pupil-sparing oculomotor nerve palsy is caused by diabetic or hypertensive oculomotor nerve ischemia, because end-arterial ischemia tends to cause more severe damage to centrally located extraoculomotor fibers than to peripherally located pupilomotor fibers [1–3]. Since oculomotor nerve palsy caused by midbrain infarction usually accompanies other neurological abnormalities [4] and oculomotor nerve palsy without other neurological findings has been considered to be rare [5], isolated pupil-sparing oculomotor nerve palsy is usually attributed to peripheral oculomotor nerve ischemia.

2. Case presentation

A 69 year-old man who took carvedilol and warfarin for hypertension and chronic atrial fibrillation experienced sudden-onset diplopia and came to our hospital 3 h after onset.

On examination, he had impaired function of the right superior, inferior, and medial rectus muscles with intact pupillary responses to light and no ptosis bilaterally. Nose–finger–nose, heel–shin, kneepat, and hand pronation and supination tests were bilaterally normal, thus not suggesting limb ataxia. He had no paralysis, extrapyramidal signs, dysarthria, or sensory disturbance. Even though his gait was apparently normal and did not include ataxic features, subtle truncal ataxia was revealed by unstable tandem gait and Mann's posture. Blood examination revealed neither diabetes mellitus nor dyslipidemia, and the prothrombin time-international normalized ratio (PT-INR) was 2.44, within therapeutic range.

Axial diffusion-weighted imaging (DWI) at 1.5-Tesla MRI showed no definite lesion; however, coronal DWI depicted a slightly hyperintense area, suggesting an acute ischemic lesion in the right para-median midbrain, which was located between the two axial sections (Fig. 1A). Two days later, although no neurological signs progressed, the lesion became more evident (Fig. 1B). MR angiography findings were within normal limits. Consequently, acute lacunar infarction was diagnosed. His diplopia improved gradually, and he had almost no visual disturbance 1 month later. The subtle truncal ataxia disappeared as well.

* Corresponding author at: 4-1-1 Otsuno, Tsuchiura-shi, Ibaraki 300-0028, Japan.

E-mail addresses: eiichiroamano1122@hotmail.co.jp (E. Amano), yamada.crad@tmd.ac.jp (I. Yamada), akinuro@tmd.ac.jp (A. Machida).




Fig. 1. An acute ischemic lesion in the right para-median midbrain with restricted diffusion on diffusion-weighted imaging (DWI) and apparent diffusion coefficient (ADC) map. Yellow arrows indicate the lesion (upper figures: axial images, each corresponding to the axial section indicated by the dotted line in the lower left-hand side figure. Lower figures: coronal images; left, DWI; right, ADC map). (A) On admission, no definite lesion was detected on axial DWI; however, coronal DWI depicted a slightly hyperintense area suggestive of acute ischemic lesion in the midbrain. The coronal ADC map also showed a slightly hypointense area. Note that the lesion is located between the axial sections. (B) Two days later, the lesion became more evident even though no neurological signs progressed during the 2 day period. The lesion is also depicted on axial DWI.

3. Discussion

In the present case, the patient showed intact pupillomotor function and no ptosis because the ischemic lesion was so small that parasympathetic fibers were spared and oculomotor fibers were not damaged completely. This case is characterized as pupil-sparing oculomotor nerve palsy without limb ataxia due to midbrain infarction, which is located below and medial to the red nucleus: a variant of Claude's syndrome that differs from the typical type in that it has no limb ataxia.

Claude's syndrome is well known as midbrain infarction causing oculomotor nerve palsy with contralateral ataxia. Seo et al [6] demonstrated that the localization of Claude's syndrome involves a lesion of the superior cerebellar peduncle (SCP) just below and medial to the red nucleus. Thereafter, ataxia in Claude's syndrome has been attributed to the destruction of the SCP, which contains the efferent fibers from the cerebellum. Tokunaga et al. [7] reported a case of midbrain infarction causing oculomotor nerve palsy with ipsilateral limb ataxia, a variant of Claude's syndrome with ipsilateral ataxia, and discussed the findings of diffusion-tensor imaging (DTI). They argued that the lesion located at the rostral extremity of the decussation of the SCP caused ipsilateral ataxia, and the destruction above the decussation resulted in the contralateral ataxia.

To identify the anatomical relationship between the ischemic lesion and the SCP, we performed DTI using a 3-Tesla MRI 27 days after the onset (Fig. 2). It revealed that the lesion was located rostrally above the decussation of the SCP and medial to the red nucleus. Because the lesion was localized adjacent to the midline and did not extend laterally and caudally, the SCP that ascends laterally after the decussation seemed to be spared. This anatomical feature explains the absence of limb ataxia in our case.

In addition, our case is characterized by the presence of subtle truncal ataxia. It is hypothesized that the ischemic lesion involved

the decussation of the tectospinal tract that locates medially to the red nucleus and the SCP [8] and that the destruction of the tectospinal tract might cause the dysfunction of postural change control. The tectospinal tract originates from the superior colliculus, decussates in the dorsal tegmental decussation medial to the red nucleus, and descends ventrolaterally to the medial longitudinal fascicle [8]. Even though the anatomy and the function of the tectospinal tract in humans largely remains undiscovered, the tectospinal tract is considered to be responsible for controlling the movement of the head and postural changes in response to the visual information [9]. Therefore, the subtle truncal ataxia in our case might be explained by the strictly localized destruction of the tectospinal tract.

In particular, in this variant of Claude's syndrome, we should carefully assess truncal ataxia to avoid misdiagnosing it as isolated pupil-sparing oculomotor nerve palsy because the patient showed apparently normal gait and truncal ataxia was only revealed by unstable tandem gait. As shown in our case, to properly diagnose pupil-sparing oculomotor nerve palsy caused by midbrain infarction, we need more careful neurological examination and additional coronal DWI because it is often difficult to detect radiologically a small midbrain infarction on axial DWI alone [10].

In summary, we have presented the case of a variant of Claude's syndrome, which shows pupil-sparing oculomotor nerve palsy without the accompanying neurological deficits other than subtle truncal ataxia. Based on the DTI, midbrain infarction was located rostrally above the decussation of the SCP and might have partially destroyed the tectospinal tract.

List of funding sources

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.




Fig. 2. Anatomical relationship of the ischemic lesion and SCP based on Diffusion-tensor imaging (DTI). Serial DWI (left) and color fractional anisotropy (FA) maps (middle) (day 27) show that the lesion (black arrow) is located rostrally above the decussation of the SCP (white arrows). Anatomical illustrations were depicted (right). RN = red nucleus, 3N = oculomotor nucleus, SC = superior colliculus, red dotted arrow = tectospinal tract, SCP = superior cerebellar peduncle, dSCP = decussation of the SCP. DTI was performed using a 3-Tesla magnetic resonance imaging (MRI) unit (Philips, Netherlands). The directions of diffusion are displayed in three colors as follows: red, right–left direction; green, anterior–posterior direction; and blue, rostral–caudal direction. A DTI sequence was performed with spin echo-based single-shot echo planar imaging (repetition time/echo time, 8829/101 ms; field of view, 230 × 230 mm; slice thickness, 2.0 mm; average, 2; matrix, 128 × 128; contiguous axial slices, 60; and voxel size, 1.8 × 1.8 × 2 mm). Images were acquired in 128 noncollinear directions ($b = 1000 \text{ s/mm}^2$) and in one acquisition without diffusion encoding ($b = 0 \text{ s/mm}^2$).

Conflicts of Interest/disclosures

The authors declare that they have no financial or other conflicts of interest in relation to this research and its population.

Acknowledgements

We wish to acknowledge the assistance of the Department of Radiology, Tsuchiura Kyodo General Hospital.

Appendix A. Supplementary data

Supplementary data associated with this article can be found, in the online version, at <https://doi.org/10.1016/j.jocn.2017.09.027>.

References

- [1] Dreyfus PM, Hakin S, Adams RD. Diabetic ophthalmoplegia: report of a case, with postmortem study and comments on vascular supply of human oculomotor nerve. *Arch Neurol Psychiatry* 1957;77:337–49.
- [2] Asbury AK, Aldredge H, Hershberg R, Fisher CM. Oculomotor palsy in diabetes mellitus: a clinico-pathological study. *Brain* 1970;93:555–66.
- [3] Weber RB, Daroff RB, Machey EA. Pathology of oculomotor nerve palsy in diabetics. *Neurology* 1970;20:835–8.
- [4] Kim JS, Kim J. Pure midbrain infarction clinical, radiologic, and pathophysiologic findings. *Neurology* 2005;64:1227–32.
- [5] Breen LA, Hopf HC, Farris BK, Gutmann L. Pupil-sparing oculomotor nerve palsy due to midbrain infarction. *Arch Neurol* 1991;48:105–6.
- [6] Seo SW, Heo JH, Lee KY, Shin WC, Chang DI, Kim SM, et al. Localization of Claude's syndrome. *Neurology* 2001;57:2304–7.
- [7] Tokunaga M, Fukunaga K, Nakanishi R, Watanabe S, Yamanaga H. Midbrain infarction causing oculomotor nerve palsy and ipsilateral cerebellar ataxia. *Intern Med* 2014;53:2143–7.
- [8] Nieuwenhuys R, Voogd J, van Huizen C. *The Human Central Nervous System*. 4th ed. Berlin Heidelberg: Springer; 2008.
- [9] Martin J. *Neuroanatomy: text and atlas*. 3rd ed. The McGraw-Hill Companies, Inc; 2003.
- [10] Oppenheim C, Stanescu R, Dormont D, Crozier S, Marro B, Samson Y, et al. False-negative diffusion-weighted MR findings in acute ischemic stroke. *Am J Neuroradiol* 2000;21:1434–40.