

Case Report

Cardiac Arrhythmias and Acute Cerebrovascular Events: A Case of QT Prolongation and Torsades de Pointes Early After Right Insular Stroke

Ilaria Jane Romano, MD, Antonio Lippolis, MD, Margherita D'Anna, MD, and Francesco Gentile, MD

Stroke involving some areas of the cerebral hemisphere, such as insula, amygdala, and lateral hypothalamus, may cause changes in autonomic control of cardiac function. A 58-year-old woman presented to the emergency department for acute onset of left facial-brachial-crural hemiparesis and dysarthria. A brain CT scan showed subacute ischemic lesion with hemorrhagic infarction in right insular-rolandic cortex. Over the next few days ECG showed severe bradycardia with elongation of QTc, significant pauses (5 seconds), runs of nonsustained ventricular tachycardia and torsades de pointes. Drug induced and other several possible causes of elongation of QT and bradycardia such as hypokalemia, a history of heart failure, and structural heart disease were ruled out. The case confirms that insular cortex plays a major role in stroke-induced cardiovascular changes.

Key Words: Ischemic stroke—right insular stroke—cardiac arrhythmias—cardiac monitoring, QTc prolongation—torsades de pointes
© 2019 Elsevier Inc. All rights reserved.

Case Report

A 58-year-old woman presented to the emergency department for acute onset of left facial-brachial-crural hemiparesis and dysarthria. Clinical history was positive for arterial hypertension, rheumatoid arthritis, and previous minor stroke. Her medications immediately before hospital admission included losartan 50 mg, and methotrexate 7.5 mg every week. Last year she underwent cardiological investigations for atypical chest pain. Serial troponin I was negative. An echocardiogram showed moderate hypertrophy of left ventricle with good systolic function and normal valves and an Holter

ECG did not show arrhythmias. ECG at admission showed new onset of atrial fibrillation (AF) with fast ventricular response. Brain computed tomography at admission showed no focal lesions; a brain computed tomography scan was again performed after 48 hours and showed subacute ischemic lesion with hemorrhagic infarction in right insular-rolandic cortex. Three days after the acute cerebrovascular accident, ECG continuous monitoring showed brady-tachy syndrome characterized by alternate AF with high ventricular response and sinus bradycardia with significant asymptomatic pauses (>2 seconds) and long QT (Fig 1). On day 4, patient presented torsades de pointes triggered by long QT with spontaneous regression (Fig 2). The day after, ECG showed severe sinus bradycardia (HR 30 bpm) with further prolongation of QTc (570 milliseconds), runs of nonsustained ventricular tachycardia and significant pauses (5 seconds). For this reason, a temporary pacemaker was first implanted and then, on day 6 after ischemic stroke, a definitive pacemaker was positioned. After definitive implantation of pacemaker, ventricular tachycardia was no longer detected at ECG continuous monitoring.

From the Cardiology Department, ASST Nord Milano, Ospedale E. Bassini, Milano, Italy.

Received January 22, 2019; revision received June 17, 2019; accepted July 21, 2019.

Address correspondence to Antonio Lippolis MD, Department of Cardiology, ASST Nord Milano-Bassini Hospital, Via M. Gorkj 50; 20092, Cinisello Balsamo, Milan, Italy. E-mail: alippolis2002@yahoo.it.
1052-3057/\$ - see front matter

© 2019 Elsevier Inc. All rights reserved.

<https://doi.org/10.1016/j.jstrokecerebrovasdis.2019.104308>




Figure 1. Sinus bradycardia with significant asymptomatic pauses (>2 seconds) and QT elongation.

Discussion

Stroke involving both cerebral hemispheres may cause changes in autonomic control of cardiac function.¹ However, some areas of the cerebral hemisphere seem to be mostly involved. In particular, the insular cortex within the middle cerebral artery territory is the most important cortical area involved in the control of both parasympathetic and sympathetic cardiovascular activity. Because middle cerebral artery occlusion is the most frequent cause of ischemic stroke, insular cortex is frequently involved in ischemic injury. Experimental animal studies demonstrated a crucial role of the insula in

stroke-induced cardiovascular and autonomic disturbances such as changes in arterial pressure, heart rate, respiration, and adrenaline secretion.⁸⁻⁹ Moreover, human studies demonstrated that stroke in the region of insula, especially the right one, lead to decreased heart rate variability and to increased incidence of sudden death.²⁻¹⁰ These findings altogether demonstrate that right insular cortex plays a major role in the autonomic control of cardiac function and suggest that a stroke localized in this area may cause a change in sympatho-vagal balance in favor of increased sympathetic tone.⁶⁻¹³ As a consequence of the increased sympathetic tone,




Figure 2. Torsades de pointes presented on day 4.

acute severe hypertension, cerebrogenic pulmonary edema caused by transient increase in pulmonary intravascular pressure and cardiac arrhythmias may occur. Cardiac arrhythmias include bradycardia, supraventricular tachycardia, atrial flutter, AF, ectopic ventricular beats, multifocal ventricular tachycardia, torsade de pointes, and ventricular fibrillation.⁵⁻⁷ Furthermore, some patients seem to be more vulnerable than others for the development of ventricular arrhythmia after stroke. In particular, a number of studies have identified possible risk factors such as older age, stroke severity with larger size of brain lesion, and reduced heart rate.^{14,15} Also, females have been shown to be more vulnerable than males in developing both hypokalemia and QT interval prolongation after acute stroke.¹⁶ In our case, after ruling out several possible causes of QT prolongation and bradycardia such as drug therapy, hypothyroidism, congenital long QT syndrome and electrolyte alterations, a neurogenic cause of arrhythmia due to insular involvement was hypothesized.⁵⁻¹⁷ This case confirms that insular cortex plays a major role in stroke-induced cardiovascular changes. Moreover, the increased risk of arrhythmias following an acute stroke in patient with newly developed ECG changes strengthens the concept that cardiac monitoring even in Neurology Department and the collaboration between Neurologist and Cardiologist is the key role to receive optimal cardiology therapy which may in turn improve quality of life and short-/long-term survival. Antiadrenergic agents that limit the effects of sympathetic response could probably be useful for the prevention or treatment of these arrhythmias. On the other hand, beta blockers should be skeptically administrated in patients with bradycardia or heart block. Finally, we believe that further studies are needed to better identified patients who carry higher risk of developing malignant arrhythmia after acute stroke and that can therefore benefit of more intensive cardiac monitoring.

Conflicts of Interest

All authors of the article confirm that there are no conflicts of interest.

References

1. Al-Qudah ZA, Yacoub HA, Souayah N. Disorders of the autonomic nervous system after hemispheric cerebrovascular disorders: an update. *J Vasc Interv Neurol* 2015;8:43-52.
2. Prosser J, MacGregor L, Lees KR, et al. VISTA Investigators. Predictors of early cardiac morbidity and mortality after ischemic stroke. *Stroke* 2007;38:2295.
3. Oppenheimer SM, Cecchetto DF, Hachinski VC. Cerebrogenic cardiac arrhythmias. Cerebral electrocardiographic influences and their role in sudden death. *Arch Neurol* 1990;47:513-519.
4. Silver FI, Norris JW, Lewis AJ, et al. Early mortality following stroke: a prospective review. *Stroke* 1984;15:492-496.
5. Cheung RT, Hachinski VC. The insula and cerebrogenic sudden death. *Arch Neurol* 2000;57:1685-1688.
6. Effects of insula resection on autonomic nervous system activity. *Neurosurg Anesthesiol* 2016;28:153-158.
7. Zini A, Fioravanti V, Ferraro D, et al. Paroxysmal ventricular tachycardia and pure right insular stroke. *J Cardiovascular Med* 2012;13:842-843.
8. Cecchetto DF, Wilson JX, Smith KE, et al. Autonomic and myocardial changes in middle cerebral artery occlusion: stroke models in the rat. *Brain Res* 1989;502:296-305.
9. Oppenheimer SM, Cecchetto DF. Cardiac chronotropic organization of the rat insular cortex. *Brain Res* 1990;533:66-72.
10. Tokgozoglul SL, Bartur MK, Topcuoglu MA, et al. Effect of stroke localization on cardiac autonomic balance and sudden death. *Stroke* 1999;30:1307-1311.
11. Oppenheimer SM, Kedem G, Martin WM. Left insular cortex lesions perturb cardiac autonomic tone in humans. *Clin Auton Res* 1996;6:131-140.
12. Sander D, Klingelhofer J. Changes of circadian blood pressure patterns after hemodynamic and thromboembolic brain infarction. *Stroke* 1994;25:1730-1737.
13. Oppneheimer SM, Gelb A, Girvin GP, et al. Cardiovascular effects of human insular cortex stimulation. *Neurology* 1992;42:1727-1732.
14. Frangiskakis JM, Hravnak M, Crago EA, et al. Ventricular arrhythmia risk after subarachnoid hemorrhage. *Neurocrit Care* 2009;10:287-294.
15. Fernández-Menéndez S, García-Santiago R, Vega-Primob A, et al. Cardiac arrhythmias in stroke unit patients. Evaluation of the cardiac monitoring data. *Neurologia* 2016;31:289-295.
16. Fukui S, Otani N, Katoh H, et al. Female gender as a risk for hypokalemia and QT prolongation after subarachnoid hemorrhage. *Neurology* 2002;59:134-136.
17. Lowattana S, Zeger SL, Lima JA, et al. Left insular stroke is associated with adverse cardiac outcome. *Neurology* 2006;66:477-483.